arXiv:1611.02331v1 [math.AG] 7 Nov 2016

NEW FAMILIES OF RANK 2 BUNDLES ON PROJECTIVE SPACE

CHARLES ALMEIDA; MARCOS JARDIM

Abstract. We present a new family of monads whose cohomology is a stable rank two vector bundle on $\mathbb{P}^3$. We also study the irreducibility and smoothness together with a geometrical description of some of these families. Such facts are used to prove that the moduli space of stable rank two vector bundles of zero first Chern class and second Chern class equal to 5 has exactly three irreducible components.

1. Introduction

In [17] Maruyama proved that the rank r stable vector bundles on a projective variety X with fixed Chern classes $c_1, ..., c_r$ can be parametrized by an algebraic quasi-projective variety, denoted by $\mathcal{B}_X(r, c_1, ..., c_r)$. Although this result has been known for almost 40 years, there are just a few concrete examples and established facts about such varieties, even for cases like $X = \mathbb{P}^3$ and $r = 2$. For instance, $\mathcal{B}_{\mathbb{P}^3}(2, 0, 1)$ was studied by Barth in [2], $\mathcal{B}_{\mathbb{P}^3}(2, 0, 2)$ was described by Harthorne in [9], $\mathcal{B}_{\mathbb{P}^3}(2, -1, 2)$ studied by Harthorne and Sols in [11] and $\mathcal{B}_{\mathbb{P}^3}(2, -1, 4)$ described by Banica and Manolache in [1]. This probably happened due to the fact that the questions of irreducibility (solved in [21] and [20]), and smoothness (answered in [16]) of the so-called *instanton component* of the moduli space $\mathcal{B}_{\mathbb{P}^3}(2, 0, c_2)$ remained opened until 2014.

In this paper, we continue the study of the moduli space $\mathcal{B}_{\mathbb{P}^3}(2, 0, n)$, which we will simply denote by $\mathcal{B}(n)$ from now on, with the goal of providing new examples of families of vector bundles, and understanding their geometry. It is more or less clear from the table in [10, Section 5.3] that $\mathcal{B}(1)$ and $\mathcal{B}(2)$ are irreducible, while $\mathcal{B}(3)$ and $\mathcal{B}(4)$ have exactly two irreducible components; the first interesting case is $\mathcal{B}(5)$. For $n \geq 6$, two families of irreducible components have been studied, namely the *instanton components*, whose generic point corresponds to an instanton bundle, and the *Ein components*, whose generic point corresponds to a bundle given as cohomology of monads of the form

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-c) \xrightarrow{\alpha} \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \oplus \mathcal{O}_{\mathbb{P}^3}(b) \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(c) \to 0$$

Date: June, 2016.

where $c > b \geq a \geq 0$ and $c > a + b$. All of the components of $\mathcal{B}(n)$ for $n \leq 4$ are of either of these types; here, we focus on certain new families of bundles that appear as soon as $n \geq 5$.

More precisely, we produce examples of families of vector bundles in $\mathcal{B}(a^2 + b^2)$, for every pair of integers a, b such that $a \geq 2b$, namely families of vector bundles arising as cohomology of monad of the form:

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} 6 \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0$$

Which will be denoted by $\mathcal{F}(a, b)$. We also provide a bijection of the family of vector bundles arising as cohomology of the monad :

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \oplus \mathcal{O}_{\mathbb{P}^3}(-1)^k \xrightarrow{\alpha} (4 + 2k) \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(a) \oplus \mathcal{O}_{\mathbb{P}^3}(1)^k \to 0$$

and the family of vector bundles arising as cohomology of

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\sigma} \tilde{E} \xrightarrow{\tau} \mathcal{O}_{\mathbb{P}^3}(a) \to 0$$

where $\tilde{E}$ is a rank 4 instanton bundle of charge k.

These facts are used to prove that $\mathcal{F}(a, 1)$ is a irreducible dense open subset of the total space of a vector bundle over $\mathcal{I}(2, 1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$ where $\mathcal{I}(2, 1)$ is the moduli space of rank two instanton bundles of second Chern class 1 and $G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$ is the grassmannian variety of planes in $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a))$. Furthermore, considering the following monads:

$$0 \to 5 \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \to 12 \cdot \mathcal{O}_{\mathbb{P}^3} \to 5 \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{1}$$

$$0 \to 2 \cdot \mathcal{O}_{\mathbb{P}^3}(-2) \to 3 \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \oplus 3 \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 2 \cdot \mathcal{O}_{\mathbb{P}^3}(2) \to 0 \tag{2}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-3) \to \mathcal{O}_{\mathbb{P}^3}(-2) \oplus 2 \cdot \mathcal{O}_{\mathbb{P}^3} \oplus \mathcal{O}_{\mathbb{P}^3}(2) \to \mathcal{O}_{\mathbb{P}^3}(3) \to 0 \tag{3}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-2) \oplus \mathcal{O}_{\mathbb{P}^3}(-1) \to 6 \cdot \mathcal{O}_{\mathbb{P}^3} \to \mathcal{O}_{\mathbb{P}^3}(1) \oplus \mathcal{O}_{\mathbb{P}^3}(2) \to 0 \tag{4}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-2) \oplus 2 \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\alpha} \mathcal{O}_{\mathbb{P}^3}(-1) \oplus 6 \cdot \mathcal{O}_{\mathbb{P}^3} \oplus \mathcal{O}_{\mathbb{P}^3}(1) \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(2) \oplus 2 \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{5}$$

we are able to prove our main result.

Main Theorem. The moduli space $\mathcal{B}(5)$ has exactly 3 irreducible components. Namely:

i) the *instanton component*, of dimension 37, which contains the bundles given as cohomology of monads of the form (1) and (2)

ii) the *Ein component*, of dimension 40, which contains the bundles given as cohomology of monads of the form (3)

iii) the closure of the family $\mathcal{F}(2,1)$, of dimension 37, which contains the bundles given as cohomology of monads of the form (4) and (5).

This paper will be organized as follows: in the Section 2 we fix the notation and conventions, in Section 3 we state the main facts about instanton sheaves that will be used, in Section 4 we construct a family of vector bundles lying in $\mathcal{B}(a^2+b^2)$, for every pair of integers a, b, such that $a \geq 2b$, and prove some facts about the dimension of this family, in Sections 5 and 6 we prove the irreducibility and smoothness of such families for the case $b = 1$, and finally, in Section 7 we study the variety $\mathcal{B}(5)$.

Acknowledgements. CA is supported by the FAPESP grant number 2014/08306-4. MJ is partially supported by the CNPq grant number 303332/2014-0 and the FAPESP grant number 2014/14743-8. Both authors would like to thank Alexander S. Tikhomirov, who shared with us his conjecture that $\mathcal{B}(5)$ has exactly three components of dimensions 37, 37 and 40; we thank him for many stimulating conversations and valuable comments during the preparation of this work. CA would also like to thank Aline Andrade for many suggestions on the presentation of the paper, and useful comments in the proof of Propositions 12 and 13.

Notation and Conventions

In this work $\mathbb{K}$ is an algebraically closed field of zero characteristic; $\mathbb{P}^3 := \mathrm{Proj}(\mathbb{K}[x,y,z,w])$. We will not make any distinction between vector bundles and locally free sheaves, and $\mathrm{H}^i(\mathcal{F})$ will denote the i-th cohomology group of the sheaf $\mathcal{F}$ on $\mathbb{P}^3$. Since we are working with rank 2 vector bundles on $\mathbb{P}^3$, and Gieseker stability is equivalent to $\mu-$stability, we will not make any distinction between these two concepts. If V is a vector space over $\mathbb{K}$, we will denote by $G(k,V)$ the grassmannian variety of k-planes in V.

2. Monads

Recall that a monad is a complex of vector bundles of the form:

$$A \xrightarrow{\alpha} B \xrightarrow{\beta} C \tag{6}$$

such that $\beta \circ \alpha = 0$, α is injective, and β is surjective. We call the sheaf $E := \ker\beta/\mathrm{Im}\alpha$ the cohomology of the monad (6). When α is locally left invertible, then E is a vector bundle.

The notion of monad is important in the study of vector bundles on $\mathbb{P}^3$ because Horrocks proved in [12] that every vector bundle on $\mathbb{P}^3$ is cohomology of a monad of the form (6) with $\mathcal{A}$, $\mathcal{B}$ and $\mathcal{C}$ being sums of line bundles.

For completeness, we include in this section some useful results about monads that will be required in this work.

The following Lemma, gives a relation between isomorphisms classes of monads and its cohomology vector bundles. A proof can be found in [18, Lemma 4.1.3].

Lemma 1. Let E and E' be, respectively, cohomology of the following monads:

$$M: \qquad A \xrightarrow{a} B \xrightarrow{b} C \tag{7}$$

$$M': \qquad A' \xrightarrow{a'} B' \xrightarrow{b'} C' \tag{8}$$

If one has that $\mathrm{Hom}(B, A') = \mathrm{Hom}(C, B') = \mathrm{H}^1(X, C^\vee \otimes A') = \mathrm{H}^1(X, B^\vee \otimes A') = \mathrm{H}^1(X, C^\vee \otimes B') = \mathrm{H}^2(X, C^\vee \otimes A') = 0$ then there exists a bijection between the set of all morphisms from E to E' and the set of all morphisms of monads from (7) to (8).

The following important Corollary will be used several times in this work, and a proof can also be found in [18, Lemma 4.1.3, Corollary 2].

Corollary 2. Consider the monad

$$M: \qquad A \xrightarrow{a} B \xrightarrow{b} C \tag{9}$$

and its dual monad:

$$M^\vee: \qquad C^\vee \xrightarrow{b^\vee} B^\vee \xrightarrow{a^\vee} C^\vee \tag{10}$$

If these two monads satisfies the hypothesis of the Lemma 1, and there exists an isomorphism $f : E \to E^\vee$, of the cohomology bundles, such that $f^\vee = -f$ then there are isomorphisms $h : C \to C^\vee$, and $q : B \to B^\vee$, such that $q^\vee = -q$, and $h \circ b = a^\vee \circ q$

We end this section with a result that gives a relation between the maps defining the monad and the extensions group of the cohomology vector bundle. It will be important in our later studies of the dimension of the family of vector bundles arising as cohomology of monads of a given form. Its proof can be found in [18, Lemma 4.1.7].

Lemma 3. Let:

$$0 \longrightarrow A \xrightarrow{a_0} B \xrightarrow{b_0} C \longrightarrow 0 \tag{11}$$

be a monad over a complex manifold X with cohomology E. For the mapping :

$$d_0 : Hom(A, B) \oplus Hom(B, C) \to Hom(A, C) \tag{12}$$

defined by $d_0(a, b) = b_0 a + b a_0$ we have that $\operatorname{coker} d_0 \simeq \mathrm{H}^2(X, EndE)$ if, and only if the following cohomology groups vanish:
$\mathrm{H}^1(X, B^* \otimes C), \mathrm{H}^1(X, A^* \otimes B), \mathrm{H}^1(X, B^* \otimes B), \mathrm{H}^1(X, C^* \otimes C), \mathrm{H}^1(X, A^* \otimes A),$
$\mathrm{H}^2(X, B^* \otimes B), \mathrm{H}^2(X, A^* \otimes A), \mathrm{H}^2(X, C^* \otimes C), \mathrm{H}^2(X, B^* \otimes A), \mathrm{H}^2(X, C^* \otimes B)1,$
$\mathrm{H}^3(X, C^* \otimes A), \mathrm{H}^3(X, B^* \otimes A), \mathrm{H}^3(X, C^* \otimes B), \mathrm{H}^4(X, C^* \otimes A).$

3. Sympletic Instanton Bundles and Sheaves

Instanton bundles are of great importance, due to their because they many remarkable properties and many applications on mathematical physics. Beside this, The Instanton component is the only known irreducible component of the moduli space $\mathcal{B}_{\mathbb{P}^3}(c)$ for every $c \in \mathbb{N}$.

In this section we will present the main results concerning instanton sheaves and vector bundles that will be used in this work. We start by recalling the definition of instanton sheaves, cf. [13, Introduction]

Definition 4. An *instanton sheaf* on $\mathbb{P}^n$ $(n \geq 2)$ is a torsion free coherent sheaf E on $\mathbb{P}^n$ with $c_1(E) = 0$ satisfying the following cohomological conditions:

(1) for $n \geq 2$, $\mathrm{H}^0(E(-1)) = \mathrm{H}^n(E(-n)) = 0$;
(2) for $n \geq 3$, $\mathrm{H}^1(E(-2)) = \mathrm{H}^{n-1}(E(1-n)) = 0$;
(3) for $n \geq 4$, $\mathrm{H}^p(E(k)) = 0$, $2 \leq p \leq n-2$ and $\forall k$;

The integer $c = \chi(E(-1))$ is called the *charge* of E. When E is locally free, we say that E is an *instanton bundle*.

Recall also that a locally free sheaf E is *sympletic* if it admits a sympletic structure, that is, there exists an isomorphism $\varphi : E \to E^\vee$, such that $\varphi^\vee = -\varphi$. A *sympletic instanton bundle* is a pair (E, φ) consisting of an instanton bundle E together with a symplectic structure φ on it. Following the proof of [5, Theorem 3.1] is possible to see that:

Theorem 5. If (E, φ) is sympletic instanton bundle on $\mathbb{P}^3$, then E is isomorphic to the cohomology of a monad of the form:

$$0 \to \mathrm{H}^2(E(-3)) \otimes \mathcal{O}_{\mathbb{P}^3}(-1) \to \mathrm{H}^1(E \otimes \Omega^1_{\mathbb{P}^3}) \otimes \mathcal{O}_{\mathbb{P}^3} \to \mathrm{H}^2(E(-3))^\vee \otimes \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{13}$$

In addition, $\mathrm{H}^1(E \otimes \Omega^1_{\mathbb{P}^3})$ admits a natural sympletic structure as a vector space.

Conversely, we observe that the existence of a sympletic structure on $\mathrm{H}^1(E \otimes \Omega^1_{\mathbb{P}^3})$ is enough to guarantee that E also admits a sympletic structure, by Corollary (2)and [14, Lemma 2.2].

The cokernel N of any monomorphism of sheaves $\mathcal{O}_{\mathbb{P}^3}(-1) \to \Omega_{\mathbb{P}^3}(1)$, is called a *null correlation sheaf*. It is straightforward to check that N is a rank 2 instanton sheaf of charge 1. If N is locally free, we say that N is a *null correlation bundle*. The following fact seems to well known, but for lack of a reference we include a proof here.

Lemma 6. Every rank 4 instanton bundle E of charge 1 over $\mathbb{P}^3$ fits into a sequence:

$$0 \to 2 \cdot \mathcal{O}_{\mathbb{P}^3} \to E \to N \to 0 \tag{14}$$

where N is a null correlation sheaf. In addition, if E admits a symplectic structure, then N is a null correlation bundle, and hence $E = N \oplus 2 \cdot \mathcal{O}_{\mathbb{P}^3}$

Proof. If E is a rank 4 instanton bundle of charge 1, then, by [6], E is 1-regular, so that $h^0(E) = \chi(E) = 2$. It follows that there exists an injective morphism of sheaves $2 \cdot \mathcal{O}_{\mathbb{P}^3} \to E$. Computing cohomologies, one sees that the cokernel of this morphism is a null correlation sheaf, and we obtain the claimed exact sequence.

Now, assuming that E admits a symplectic structure, it is enough to show that $\mathrm{H}^1(N \otimes \Omega^1_{\mathbb{P}^3})$ admits a natural functorial sympletic structure. Twisting the sequence (14) by $\Omega^1_{\mathbb{P}^3}$ and $\Omega^2_{\mathbb{P}^3}$ we obtain the following two exact sequences of cohomology, respectively:

$$0 \to 2 \cdot \mathrm{H}^1(\Omega^1_{\mathbb{P}^3}) \to \mathrm{H}^1(E \otimes \Omega^1_{\mathbb{P}^3}) \to \mathrm{H}^1(N \otimes \Omega^1_{\mathbb{P}^3}) \to 0 \tag{15}$$

$$0 \to 2 \cdot \mathrm{H}^2(\Omega^2_{\mathbb{P}^3}) \to \mathrm{H}^2(E \otimes \Omega^2_{\mathbb{P}^3}) \to \mathrm{H}^2(N \otimes \Omega^2_{\mathbb{P}^3}) \to 0 \tag{16}$$

The vector space $\mathrm{H}^1(\Omega_{\mathbb{P}^3})^\vee$ admits a sympletic structure, and so does $\mathrm{H}^1(E\otimes\Omega_{\mathbb{P}^3})^\vee$ by hypothesis; they are naturally isomorphic to $\mathrm{H}^2(\Omega^2_{\mathbb{P}^3})$ and $\mathrm{H}^2(E\otimes\Omega^2_{\mathbb{P}^3})$, respectively, via Serre duality, and hence admit natural functorial sympletic structures. By comparing the duals of sequences (15) and (16), it follows that $\mathrm{H}^1(N\otimes\Omega_{\mathbb{P}^3})^\vee$ is naturally isomorphic to $\mathrm{H}^2(N\otimes\Omega^2_{\mathbb{P}^3})$, and that the isomorphism is symplectic, thus N is locally free. Since $\mathrm{Ext}^1(N,\mathcal{O}_{\mathbb{P}^3})=\mathrm{H}^1(N)=0$, the sequence (14) splits, hence $E=N\oplus 2\cdot\mathcal{O}_{\mathbb{P}^3}$. □

4. Locally free sheaves with two generators in the first cohomology group

Recall that every locally free sheaf E on $\mathbb{P}^3$ is cohomology of a monad of the form [12]:

$$0\to\oplus_{i=1}^r\mathcal{O}_{\mathbb{P}^3}(a_i)\to\oplus_{j=1}^s\mathcal{O}_{\mathbb{P}^3}(b_j)\to\oplus_{k=1}^t\mathcal{O}_{\mathbb{P}^3}(c_k)\to 0 \tag{17}$$

In this work, we will be interested in rank 2 locally free sheaves with vanishing first Chern class. Under these conditions, we have $E^\vee\simeq E$, thus the monad (17) is self dual, which implies $t=r$, $s=2r+2$, and $\{a_i\}=-\{c_k\}$. In addition, the middle entry of the monad is also self dual, so that (17) reduces to

$$0\to\oplus_{i=1}^r\mathcal{O}_{\mathbb{P}^3}(a_i)\to\oplus_{j=1}^{r+1}\left(\mathcal{O}_{\mathbb{P}^3}(b_j)\oplus\mathcal{O}_{\mathbb{P}^3}(-b_j)\right))\to\oplus_{i=1}^r\mathcal{O}_{\mathbb{P}^3}(-a_i)\to 0. \tag{18}$$

Finally, recall also that r coincides with the number of generators of $\mathrm{H}^1_*(E)=\bigoplus_{p\in\mathbb{Z}}\mathrm{H}^1(E(p))$ as a module over the ring of homogeneous polynomials on four variables, while a_i are the degrees of these generators, cf. [15, Thm. 2.3].

Ein studied in [7] the case $r=1$, introducing the so called *generalized null correlation bundles*, given as cohomologies of monads of the form

$$0\to\mathcal{O}_{\mathbb{P}^3}(-a)\to\oplus_{j=1}^2\left(\mathcal{O}_{\mathbb{P}^3}(b_j)\oplus\mathcal{O}_{\mathbb{P}^3}(-b_j)\right))\to\mathcal{O}_{\mathbb{P}^3}(a)\to 0. \tag{19}$$

In addition, the cohomology of such a monad is stable if and only if $a>b_1+b_2$.

Our goal now is to study stable rank 2 locally free sheaves E with vanishing first Chern class whose cohomology module $\mathrm{H}^1_*(E)$ has exactly two generators. We begin with the following statement.

Lemma 7. A rank 2 locally free sheaf E on $\mathbb{P}^3$ is the cohomology of a monad of the form

$$0\to\mathcal{O}_{\mathbb{P}^3}(-b)\oplus\mathcal{O}_{\mathbb{P}^3}(-a)\xrightarrow{\alpha}6\cdot\mathcal{O}_{\mathbb{P}^3}\xrightarrow{\beta}\mathcal{O}_{\mathbb{P}^3}(b)\oplus\mathcal{O}_{\mathbb{P}^3}(a)\to 0 \tag{20}$$

if and only if $c_1(E)=0$, $c_2(E)=a^2+b^2$ and $\mathrm{H}^1_*(E)$ has exactly two generators in degrees $-a$ and $-b$.

Proof. The first part is straightforward. If E is a rank 2 locally free sheaf with vanishing first Chern class such that $\mathrm{H}^1_*(E)$ is generated in degrees $-a$ and $-b$, then by [15, Theorem 2.3], E is cohomology of a monad of the form:

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} \oplus_{i=1}^{6} \mathcal{O}_{\mathbb{P}^3}(k_i) \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0.$$

A simple calculation gives us that $c_2(E) = a^2 + b^2 - \sum_{i=1}^{6} k_i^2$; since $c_2(E) = a^2 + b^2$, we have $k_i = 0$ for all i, as desired. □

Barth [3] and Hartshorne and Rao [10] have considered monads of the form (20) for some values of a and b, particularly $a = 2$ and $b = 1$. More generally, we start by checking that monads of the form (20) do exist for any pair of integers a and b. Indeed, assume $a \geq b$, and consider the following morphisms:

$$\alpha = \begin{pmatrix} y^a & w^b \\ -x^a & -z^b \\ -z^b y^{a-b} & 0 \\ w^b x^{a-b} & 0 \\ z^a & y^b \\ -w^a & -x^b \end{pmatrix}, \tag{21}$$

$$\beta = \begin{pmatrix} x^a & y^a & z^a & w^a & z^b y^{a-b} & w^b x^{a-b} \\ z^b & w^b & y^b & x^b & 0 & 0 \end{pmatrix}. \tag{22}$$

One easily checks that $\beta \circ \alpha = 0$; moreover, note that the following minors of both matrices

$$y^{a+b} - w^b z^a \ , \ x^a y^b - z^{a+b} \ , \ x^{a+b} - z^b w^a$$

$$y^a x^b - w^{a+b} \text{ and } x^b y^{a-b} z^b$$

do not vanish simultaneously. It follows that the cohomology sheaf is locally free.
The next problem is to establish the stability of the vector bundles arising as cohomology of monads of the form (20), depending on a and b. The following lemma is a technical result that help us to obtain a partial answer to this problem.

Lemma 8. If a, b are positive integers such that $a \geq 2b > 0$, and α and β are morphisms of vector bundles, then does not exist a monad of the type below:

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} 5 \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{23}$$

Proof. Supose, by contradiction, that there exists such monad, and let $K = \ker \beta$ and E the vector bundle that is cohomology of this monad, we have the following exacts sequences of vector bundles :

$$0 \to K \to 5 \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{24}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} K \to E \to 0 \tag{25}$$

Dualizing 24 we get :

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} 5 \cdot \mathcal{O}_{\mathbb{P}^3} \to K^\vee \to 0 \tag{26}$$

By [4, Theorem 2.7] $K^\vee$ is stable, and so is K. By an elementary calculation we check that the slope $\mu(K) = \frac{-b-a}{3}$, but by the injectivity of α, $\mathcal{O}_{\mathbb{P}^3}(-a)$ is a torsion free subsheaf of K with slope $\mu(\mathcal{O}_{\mathbb{P}^3}(-a)) = -a$ which can not occur because $a \geq 2b$. Hence such monads do not exist.

□

With this Lemma we are able to prove the main result of this section.

Theorem 9. Let α, β be morphisms of vector bundles, and $2a \geq b > 0$ positive integers. If E is the rank 2 bundle on $\mathbb{P}^3$ arising from the cohomology of a monad of the type :

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} 6 \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{27}$$

Then E is a self dual stable vector bundle. Furthermore, E is cohomology of (27) if, and only if $\mathrm{H}^1_*(E)$ is generated by exactly two elements in degree $-a$ and $-b$, and $c_2(E) = a^2 + b^2$

Proof. Let $K = \ker \beta$, the display of the monad gives:

$$0 \to K \xrightarrow{i} 6 \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{28}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\alpha} K \to E \to 0 \tag{29}$$

One sees that $c_1(E) = 0$ which implies, since E is rank 2 bundle, that E is self dual.

By Hoppe's Criteria E is stable if, and only if $\mathrm{H}^0(E) = 0$.

Again from the display of the monad, $\mathrm{H}^0(K) \simeq \mathrm{H}^0(E)$. Suppose, by contradiction, that $\mathrm{H}^0(K) \neq 0$, so there is a non zero global section $s \in \mathrm{H}^0(K)$ that induces a morphism $0 \to \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\cdot s} K$ the injectivity of this map follows from the fact that every morphism of a line bundle to another vector bundle, is either the zero map, or injective. The composition of this map with the inclusion on 28 gives an injective map $\psi = s \circ i : \mathcal{O}_{\mathbb{P}^3} \to 6 \cdot \mathcal{O}_{\mathbb{P}^3}$, which induces a injective map of complexes $N^\bullet \to M^\bullet$, since $\beta \circ s \circ i = 0$ because $M^\bullet$ is a monad.

(30)

$$\begin{array}{lccccccccc}
N^\bullet: & 0 & \longrightarrow & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3} & \longrightarrow & 0 & \longrightarrow & 0 \\
{\scriptstyle\Psi}\downarrow & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
M^\bullet: & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) & \xrightarrow{\alpha} & 6 \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\beta} & \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \\
\downarrow & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\operatorname{coker}\Psi: & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-b) \oplus \mathcal{O}_{\mathbb{P}^3}(-a) & \xrightarrow{\alpha'} & 5 \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\beta'} & \mathcal{O}_{\mathbb{P}^3}(b) \oplus \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0
\end{array}$$

Therefore coker Ψ is a monad. But it can not occur by Lemma 8. Then $\mathrm{H}^0(E) \simeq \mathrm{H}^0(K) = 0$ hence E is stable. The second claim follows from Proposition 7. □

Theorem (9), shows that, for $a \geq 2b$, there is a family of rank 2 vector bundles lying on $\mathcal{B}_{\mathbb{P}^3}(2, 0, a^2 + b^2)$, arising from the cohomology of these monads. We will denote the set of the isomorphisms classes of such bundles by $\mathcal{F}(a, b)$.

The next task will be to find the dimension of this family. The following result due to Barth will be important; its proof can be found in [3, Section 5].

Proposition 10. Let $0 \to A^\vee \to B \to A \to 0$ be a self-dual monad, where the arrows indicates maps of vector bundles, and A, B are direct sum of line bundles on $\mathbb{P}^3$. Supose that every vector cohomology of such monad is stable, and that $Hom(A, B) = 0$, then the dimension of the family of isomorphisms classes of bundles E arising from this monads is:

$$\dim Hom(B, A) - \mathrm{H}^0(\Lambda^2 A) - \dim Gl(A) - \mathrm{H}^0(S^2 B) \tag{31}$$

Corollary 11. Let $a \geq 2b$, then $\mathcal{F}(a, b)$ is a locally closed subscheme of $\mathcal{B}_{\mathbb{P}^3}(2, 0, a^2 + b^2)$ of dimension

$$6\binom{a+3}{3}+6\binom{b+3}{3}-\binom{a+b+3}{3}-\binom{b-a+3}{3}-23 \tag{32}$$

It is noteworthy that for $a=2, b=1$, the family of vector bundles arising as cohomology of the monad (20) has dimension 37, which coincides with the expected dimension of an irreducible component of $\mathcal{B}(5)$. This family will be further discussed in the next sections.

5. The relation with instanton bundles

Our main goal in this section is to provide a relation between the monads that defines the vector bundles that we are interested and monads that have instanton bundles in the central position. It will help us to give a geometrical to the family $\mathcal{F}(a,b)$ for some values of a and b.

In order of investigate the geometrical properties of $\mathcal{F}(a,b)$, will be introduced the monad below.

$$0\to\mathcal{O}_{\mathbb{P}^3}(-a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\xrightarrow{\alpha}(4+2k)\cdot\mathcal{O}_{\mathbb{P}^3}\xrightarrow{\beta}\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1)\to 0 \tag{33}$$

The family of isomorphisms classes of vector bundles arising as cohomology of such monads will be denoted by $\mathcal{G}(a,k)$. Note that by now, $\mathcal{G}(a,k)$ could possible be empty. To see that it can not happen for $a\geq k$, one needs the following two results.

Proposition 12. Given a vector bundle $E\in\mathcal{G}(a,k)$ there exists a rank 4 instanton bundle $\tilde{E}$ of charge k, and morphisms σ and τ such that:

$$0\to\mathcal{O}_{\mathbb{P}^3}(-a)\xrightarrow{\sigma}\tilde{E}\xrightarrow{\tau}\mathcal{O}_{\mathbb{P}^3}(a)\to 0 \tag{34}$$

is a monad whose cohomology is isomorphic to E.

Proof. Define $\tilde{\alpha}=\alpha\circ i$ and $\tilde{\beta}=\pi\circ\beta$ where $i: k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\to\mathcal{O}_{\mathbb{P}^3}(-a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)$ is the inclusion and $\pi:\mathcal{O}_{\mathbb{P}^3}(-a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\to k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)$ is the projection. It is clear that $\tilde{\alpha}$ is injective and $\tilde{\beta}$ is surjective.

Then we get the following monad whose cohomology is a rank 4 instanton $\tilde{E}$ of charge k:

$$0\to k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\xrightarrow{\tilde{\alpha}}(4+2k)\cdot\mathcal{O}_{\mathbb{P}^3}\xrightarrow{\tilde{\beta}}k\cdot\mathcal{O}_{\mathbb{P}^3}(1)\to 0 \tag{35}$$

Now we need to construct the morphisms σ and τ.

It is straightfoward to check that the chain of inclusions: $\mathrm{Im}\tilde{\alpha} \subseteq \mathrm{Im}\alpha \subseteq \ker\beta \subseteq \ker\tilde{\beta}$ holds. For this reason we have:

(36)
$$\begin{array}{ccccccccc}
0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \ker\beta & \longrightarrow & (4+2k)\cdot\mathcal{O}_{\mathbb{P}^3} & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle f_1} & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \ker\tilde{\beta} & \xrightarrow{i} & (4+2k)\cdot\mathcal{O}_{\mathbb{P}^3} & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & \longrightarrow & \mathrm{coker}\, f_1 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0
\end{array}$$

where f_1 is the inclusion map.

And then we see that $\mathrm{coker}\, f_1 \simeq \mathcal{O}_{\mathbb{P}^3}(a)$. And we have (where ω is the inclusion):

(37)
$$\begin{array}{ccccccccc}
0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & \mathrm{Im}\tilde{\alpha} & \xrightarrow{i} & \ker\beta & \longrightarrow & \ker\beta/\mathrm{Im}\tilde{\alpha} & \longrightarrow & 0 \\
 & & \| & & \downarrow{\scriptstyle f_1} & & \downarrow{\scriptstyle \omega} & & \\
0 & \longrightarrow & \mathrm{Im}\tilde{\alpha} & \xrightarrow{i} & \ker\tilde{\beta} & \longrightarrow & \tilde{E} & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow{\scriptstyle \tau} & & \\
 & \longrightarrow & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}_3}(a) & \longrightarrow & \mathcal{O}_{\mathbb{P}_3}(a) & \longrightarrow & 0
\end{array}$$

Now, by the isomorphism theorem we have $\dfrac{\ker\beta}{\mathrm{Im}\alpha} \simeq \dfrac{\frac{\ker\beta}{\mathrm{Im}\tilde{\alpha}}}{\frac{Im\alpha}{\mathrm{Im}\tilde{\alpha}}}$, so the map f_2 is well defined, and the bellow diagram commutes:

(38)
$$\begin{array}{ccccccccc}
0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & \ker f_2 & \longrightarrow & \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}_3}(-1) & \xrightarrow{\tilde{\alpha}} & \ker\beta & \longrightarrow & \ker\beta/\mathrm{Im}\tilde{\alpha} & \longrightarrow & 0 \\
 & & \downarrow & & \| & & \downarrow{\scriptstyle f_2} & & \\
0 & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}_3}(-1)\oplus\mathcal{O}_{\mathbb{P}_3}(-a) & \longrightarrow & \ker\beta & \longrightarrow & E & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
 & \longrightarrow & \mathcal{O}_{\mathbb{P}_3}(-a) & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0
\end{array}$$

And then $\ker f_2 \simeq \mathcal{O}_{\mathbb{P}_3}(-a)$. Hence we got the exact sequence:

$$0 \to \mathcal{O}_{\mathbb{P}_3}(-a) \xrightarrow{\sigma'} \ker\beta/\mathrm{Im}\tilde{\alpha} \xrightarrow{f_2} E \to 0 \tag{39}$$

Since σ' and ω are injective, we have that $\sigma = \omega \circ \sigma'$ is a map $\sigma : \mathcal{O}_{\mathbb{P}_3}(-a) \to \tilde{E}$.
And then we have constructed the monad:

$$0 \to \mathcal{O}_{\mathbb{P}_3}(-a) \xrightarrow{\sigma} \tilde{E} \xrightarrow{\tau} \mathcal{O}_{\mathbb{P}_3}(a) \to 0 \tag{40}$$

such that E is its cohomology. □

Proposition 13. If E is cohomology of a monad of the form (34) then E is isomorphic to a vector bundle cohomology of a monad of the form (33), thus $E \in \mathcal{G}(a, k)$.

Proof. Now, let $\tilde{E}$ be an rank 4 instanton of charge k over $\mathbb{P}^3$, then $\tilde{E}$ is cohomology of a monad of the type:

$$0 \to k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\tilde{\alpha}} (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\tilde{\beta}} k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0$$

Let $\tau, \sigma \in \mathrm{H}^0(\tilde{E}(a))$, and $\tau \circ \sigma = 0$. So we have the following exacts sequences:

$$0 \to \ker\tilde{\beta} \to (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\tilde{\beta}} k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{41}$$

$$0 \to k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\tilde{\alpha}} \ker\tilde{\beta} \to \tilde{E} \to 0 \tag{42}$$

$$0 \to \ker\tau \to \tilde{E} \xrightarrow{\tau} \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{43}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\sigma} \ker\tau \to E \to 0 \tag{44}$$

where $E \simeq \ker\tau/\mathrm{Im}\sigma$
We have the trivial extension:

$$0 \to k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \to \ker\tau \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{f_1} \ker\tau \to 0 \tag{45}$$

Fix once and for all, each representative class of the elements of $\tilde{E} = \ker\tilde{\beta}/\mathrm{Im}\tilde{\alpha}$ and define a map $f_2 : \ker\tau \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \to \ker\tilde{\beta}$ such that $f_2(\overline{x}, y) = x + \tilde{\alpha}y$. Note that there

exists a family of maps of this type, one map for each choice of representatives, but by the diagram below, we see that the kernel and cokernel of these maps are always isomorphic by the snake lemma.

$$\begin{array}{ccccccccc} (46) & 0 & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}^3}(-1) & \longrightarrow & \ker\tau\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1) & \longrightarrow & \ker\tau & \longrightarrow 0 \\ & & & \downarrow & & \downarrow{\scriptstyle f_2} & & \downarrow & \\ & 0 & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}^3}(-1) & \xrightarrow{\tilde{\alpha}} & \ker\tilde{\beta} & \longrightarrow & \tilde{E} & \longrightarrow 0 \end{array}$$

And again using the Snake Lemma in the above diagram we get the exact sequence:

$$0\to\ker\tau\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\xrightarrow{f_2}\ker\tilde{\beta}\to\mathcal{O}_{\mathbb{P}^3}(a)\to 0 \tag{47}$$

Since $\ker\beta\subseteq\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1)$, f_2 can be viewed as a map, denoted by $\tilde{f}_2$ to $\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1)$, and finally, we have the commutative diagram, where the third line is obtained by the snake lemma :

$$\begin{array}{ccccccccc} (48) & 0 & \longrightarrow & \ker\tau\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1) & \longrightarrow & \ker\tau\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1) & \longrightarrow & 0 & \longrightarrow 0 \\ & & & \downarrow{\scriptstyle f_2} & & \downarrow{\scriptstyle \tilde{f}_2} & & \downarrow & \\ & 0 & \longrightarrow & \ker\tilde{\beta} & \xrightarrow{i} & (4+2k)\cdot\mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\tilde{\beta}} & k\cdot\mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow 0 \\ & & & \downarrow & & \downarrow{\scriptstyle \pi} & & \| & \\ & 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & \operatorname{coker}\tilde{f}_2 & \longrightarrow & k\cdot\mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow 0 \\ & & & \downarrow & & \downarrow & & \downarrow & \\ & & & 0 & & 0 & & 0 & \end{array}$$

Since $\operatorname{Ext}^1(k\cdot\mathcal{O}_{\mathbb{P}^3}(1),\mathcal{O}_{\mathbb{P}^3}(a))=0$, on has $\operatorname{coker}\tilde{f}_2\simeq\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1)$
Denoting $\beta=\pi$, we have:

$$0\to\ker\tau\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\xrightarrow{f_2}(4+2k)\cdot\mathcal{O}_{\mathbb{P}^3}\xrightarrow{\beta}\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1)\to 0 \tag{49}$$

Making $\alpha=f_2\circ(\sigma,id)$ we have the monad:

$$\mathcal{O}_{\mathbb{P}^3}(-a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(-1)\xrightarrow{\alpha}(4+2k)\cdot\mathcal{O}_{\mathbb{P}^3}\xrightarrow{\beta}\mathcal{O}_{\mathbb{P}^3}(a)\oplus k\cdot\mathcal{O}_{\mathbb{P}^3}(1) \tag{50}$$

whose cohomology is isomorphic to E.

□

We will denote the isomorphisms classes of vector bundles arising as cohomology of the monad (34) by $\mathcal{N}(a,k)$. The aim of the next lemma is to show that if $\tilde{E}$ is a rank 4 instanton bundle of charge k, that fits into a monad of the form (34) such that the cohomology is a vector bundle, then $\tilde{E}$ is a sympletic instanton bundle.

Lemma 14. If $\tilde{E}$ is a rank 4 instanton bundle of charge k that fits in a monad of the form (34), such that the cohomology is a vector bundle, then $\tilde{E}$ is a sympletic, and σ is determined by τ.

Proof. Since E is a rank 2 self dual vector bundle on $\mathbb{P}^3$, there is a sympletic isomorphism $\phi : E \to E^\vee$. By the Corollary 2, there is an isomorphisms of monads:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \xrightarrow{\sigma} & \tilde{E} & \xrightarrow{\tau} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \\ & & \downarrow g & & \downarrow q & & \downarrow h & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \xrightarrow{\tau^\vee} & \tilde{E}^\vee & \xrightarrow{\sigma^\vee} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \end{array} \tag{51}$$

Such that $q^\vee = -q$, hence $\tilde{E}$ is a sympletic instanton bundle, and $\sigma = (q^{-1})^\vee \circ \tau^\vee \circ h^\vee$. □

Lemma 15. For every sympletic instanton $\tilde{E}$ of rank 4 and charge k, with $a \geq k$, $\dim \mathrm{H}^0(\tilde{E}(a)) = (4+2k)\binom{a+3}{3} - k\binom{a+4}{3} - k\binom{a+2}{3}$.

Proof. By [6, Corollary 3.3], $\tilde{E}$ is k-regular, since $a \geq k$, $\tilde{E}$ is a-regular, and therefore $\mathrm{H}^1(\tilde{E}(a)) = 0$. It follows that $\chi(\tilde{E}(a)) = h^0(\tilde{E}(a)) = (4+2k)\binom{a+3}{3} - k\binom{a+4}{3} - k\binom{a+2}{3}$. □

In order to see that $\mathcal{N}(a,k)$ is non-empty for $a \geq k$, note that, by the Lemma 15, $\mathrm{H}^0(E(a))$ is not empty, hence there exists a surjective map $\tau : \tilde{E} \to \mathcal{O}_{\mathbb{P}^3}(a)$, from which we obtain a monad of the form (34). Moreover, from the definitions, one sees that $\mathcal{F}(a,1) = \mathcal{N}(a,1)$. Putting all these results together we can state the following Proposition.

Proposition 16. A vector bundle E is cohomology bundle of :

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\alpha} (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{52}$$

If, and only if ,is cohomology bundle of :

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \xrightarrow{\sigma} \tilde{E} \xrightarrow{\tau} \mathcal{O}_{\mathbb{P}^3}(a) \to 0 \tag{53}$$

Where $\tilde{E}$ is a rank 4 sympletic instanton bundle of charge k on $\mathbb{P}^3$.

An interesting cohomological characterization for the vector bundles arising from monads of the form (33) can be found in the following proposition.

Proposition 17. A vector bundle E is cohomology of the monad (33) if, and only if, $\mathrm{H}^1_*(E)$ has 1 generator in degree $-a$ and k generators in degree -1 with Chern classes $c_1(E) = 0, c_2(E) = a^2 + k$.

Proof. The "only if" part is straightforward. If E is a self dual vector bundle on $\mathbb{P}^3$ with two generators for $\mathrm{H}^1_*(E)$ in degree $-a$ and $-b$ then by [15, Theoremm 2.3], E is cohomology of a monad of the type:

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\alpha} \oplus_{i=1}^{2k+4} \mathcal{O}_{\mathbb{P}^3}(k_i) \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{54}$$

Computing the Chern class give us $c_2(E) = a^2 + k - \sum_{i=1}^{6} k_{i_1}^2$, since $c_2(E) = a^2 + b^2$, we have $k_i = 0$ for all i. Hence the result follows. □

Another question that naturally arises is if all vector bundles cohomolgy of (33) are stable. We have a partial answer to this question.

Proposition 18. There exists at least one $E \in \mathcal{G}(a, k)$ stable, for each $a \geq 1, k \in \mathbb{N}$.

Proof. Let F be an rank 2 instanton bundle of charge k. Let $a \geq 1$ and take $\sigma \in \mathrm{H}^0(F(2a))$ and $(\sigma)_0 = X$ (this section always exists if F is a 'tHooft instanton, for instance). Let Y be a complete intersection curve, of two surfaces of degree a and $X \cap Y = \emptyset$. According to [10, Lemma 4.8], there exists a bundle E and a section $\tau \in \mathrm{H}^0(E(a))$ such that $(\tau)_0 = Y \cup X$ and cohomology of a monad of the form (33). In addition, since F is stable, X is not contained in any surface of degree a, hence neither $Y \cup X$, therefore E is stable. □

The Proposition 16 tell us that $\mathcal{G}(a, k) = \mathcal{N}(a, k)$ for all $a, k \in \mathbb{N}$, and $a \geq k$ and that $\mathcal{G}(a, k)$ is not empty. The next task will be understand the isomorphisms of monads induced by the isomorphisms of vector bundles on $\mathcal{N}(a, k)$.
In order to fix the notation, note that every isomorphism $f \in \mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1), \mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1))$ can be represented by a $(k+1) \times (k+1)$ matrix :

$$f = \begin{bmatrix} f_{1,1} & 0 & \cdots & 0 \\ f_{2,1} & f_{2,2} & \cdots & f_{2,k+1} \\ \vdots & \vdots & \ddots & \vdots \\ f_{k+1,1} & f_{k+1,2} & \cdots & f_{k+1,k+1} \end{bmatrix} \tag{55}$$

where each $f_{j,1} \in \mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(-a), \mathcal{O}_{\mathbb{P}^3}(-1))$ with $j = 1...(k+1)$ and $f_{l,m}$ are constants for $l = 1, 2, ..., k+1$ and $m = 2, 3, \cdots, (k+1)$ such that:

$$f_{11}. \begin{vmatrix} f_{2,2} & \cdots & f_{2,k+1} \\ \vdots & \ddots & \vdots \\ f_{k+1,2} & \cdots & f_{k+1,k+1} \end{vmatrix} \neq 0 \tag{56}$$

we will denote :

$$\tilde{f} = \begin{bmatrix} f_{2,2} & \cdots & f_{2,k+1} \\ \vdots & \ddots & \vdots \\ f_{k+1,2} & \cdots & f_{k+1,k+1} \end{bmatrix} \tag{57}$$

clearly $\tilde{f} \in \mathrm{Aut}(k \cdot \mathcal{O}_{\mathbb{P}^3}(-1))$. And for $h \in \mathrm{Hom}(k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \oplus \mathcal{O}_{\mathbb{P}^3}(a), k \cdot \mathcal{O}_{\mathbb{P}^3}(1) \oplus \mathcal{O}_{\mathbb{P}^3}(a))$ can be represented by a $(k+1) \times (k+1)$ matrix :

$$h = \begin{bmatrix} h_{1,1} & \cdots & h_{1,k} & 0 \\ h_{2,1} & \cdots & h_{2,k} & 0 \\ \vdots & \vdots & \ddots & \vdots \\ h_{k+1,1} & h_{k+1,2} & \cdots & h_{k+1,k+1} \end{bmatrix} \tag{58}$$

where each $h_{k+1,j} \in \mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(1), \mathcal{O}_{\mathbb{P}^3}(a))$ for $j = 1, \cdots k+1$ and $h_{l,m}$ are constants for $l, m = 1, 2, \cdots, k$, such that:

$$h_{k+1,k+1}. \begin{vmatrix} h_{1,1} & \cdots & h_{1,k} \\ \vdots & \ddots & \vdots \\ h_{k,1} & \cdots & h_{k,k} \end{vmatrix} \neq 0 \tag{59}$$

We will denote:

$$\tilde{h} = \begin{vmatrix} h_{1,1} & \cdots & h_{1,k} \\ \vdots & \ddots & \vdots \\ h_{k,1} & \cdots & h_{k,k} \end{vmatrix} \tag{60}$$

clearly $\tilde{h} \in \mathrm{Aut}(k \cdot \mathcal{O}_{\mathbb{P}^3}(1))$. With this we show that:

Lemma 19. Let E and $E' \in \mathcal{N}(a,k)$. If there exists an isomorphism $G : E \to E'$ then there exists an isomorphism $g : \tilde{E} \to \tilde{E}'$ and a $\lambda \in \mathbb{K}$, such that $\tau' \circ g = \lambda\tau$ where $(\tilde{E}, \tau)$ and $(\tilde{E}', \tau')$ are the pairs associated with E and E' respectively.

Proof. Since E is rank 2 vector bundle on $\mathbb{P}^3$ with zero first Chern class, E is a sympletic vector bundle, and G is a sympletic isomorphism, by Corollary 2, , there is an skew symmetric isomorphism of monads:

(61)

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) & \xrightarrow{\alpha} & (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\beta} & \mathcal{O}_{\mathbb{P}^3}(a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \\ & & \downarrow f & & \downarrow g & & \downarrow h & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) & \xrightarrow{\alpha'} & (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\beta'} & \mathcal{O}_{\mathbb{P}^3}(a) \oplus k \cdot \mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \end{array}$$

One sees that the following diagram commutes:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & k \cdot \mathcal{O}_{\mathbb{P}^3}(-1) & \xrightarrow{\tilde{\alpha}} & (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\tilde{\beta}} & k \cdot \mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \\ & & \downarrow \tilde{f} & & \downarrow g & & \downarrow \tilde{h} & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-1)^k & \xrightarrow{\tilde{\alpha}'} & (4+2k) \cdot \mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\tilde{\beta}'} & k \cdot \mathcal{O}_{\mathbb{P}^3}(1) & \longrightarrow & 0 \end{array} \tag{62}$$

And since $\tilde{f}, g, \tilde{h}$ are isomorphisms, it follows that (62) is a isomorphism of monads. Therefore $\tilde{E} \simeq \tilde{E}'$.

From the fact that $E \simeq E'$, and that the monad given by the pair $(\tilde{E}, \tau)$ satisfies the conditions of the of Corollary 2, follows that there exists an isomorphisms of monads:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \longrightarrow & \tilde{E} & \xrightarrow{\tau} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow g & & \downarrow \lambda & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \longrightarrow & \tilde{E}' & \xrightarrow{\tau'} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \end{array} \tag{63}$$

And hence $\tau' \circ g = \lambda\tau$.

□

Let $P(a,k)$ be the set of pairs $(\tilde{E},\tau)$ with $\tilde{E}$ sympletic instanton of rank 4 and charge k, with $\tau \in \mathrm{H}^0(\tilde{E}(a))$, equipped with the following equivalence relation: $(\tilde{E},\tau) \sim (\tilde{E}',\tau')$ if, and only if, $\tilde{E} \simeq \tilde{E}'$ and there exists a non-zero constant λ such that $\tau' \circ g = \lambda\tau$ we will denote each equivalence class in $P(a,k)$ by $[\tilde{E},\tau]$.
The Lemma 19 together with the Theorem 16 give us the following proposition.

Proposition 20. There exists a bijection between the isomorphisms classes of vector bundles on $\mathcal{N}(a,k)$ and the set of equivalences classes in $P(a,k)$. Furthermore, there exists a bijection between $P(a,1)$ and $\mathcal{F}(a,1)$.

From now on we will focus on the set $\mathcal{P}(a,1)$, with $a \geq 2$. Note that when $a = 1$, by the previous Proposition, there exists a bijection between $P(1,1)$ and the family of instanton bundles on $\mathbb{P}^3$ with charge 5.

6. The structure of $P(a,1)$

In this section, our aim is to show that $P(a,1)$ is a irreducible smooth variety. We will not make any distinction between a vector bundle E and its class of isomorphisms $[E]$ and we will denote both of them by the same letter E without the brackets.
Given $[\tilde{E},\tau] \in P(a,1)$, one has that $\tilde{E}$ is a rank 4 instanton bundle of charge 1, and by Lemma 6, there exists a null correlation bundle N over $\mathbb{P}^3$ such that $\tilde{E} \simeq N \oplus 2\cdot\mathcal{O}_{\mathbb{P}^3}$. Now, if $(\tilde{E},\tau)$ and $(\tilde{E}',\tau')$ are in the same equivalence class $[\tilde{E},\tau]$, then $\tilde{E} \simeq \tilde{E}' \simeq N \oplus 2\cdot\mathcal{O}_{\mathbb{P}^3}$ and we have that the following commuative diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \longrightarrow & N \oplus 2\cdot\mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\tau} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \\ & & \downarrow & & \downarrow g & & \downarrow \lambda & & \\ 0 & \longrightarrow & \mathcal{O}_{\mathbb{P}^3}(-a) & \longrightarrow & N \oplus 2\cdot\mathcal{O}_{\mathbb{P}^3} & \xrightarrow{\tau'} & \mathcal{O}_{\mathbb{P}^3}(a) & \longrightarrow & 0 \end{array} \tag{64}$$

Note that, since N is stable, that g can be represented by :

$$g = \begin{pmatrix} 1 & 0 & 0 \\ 0 & a_1 & a_2 \\ 0 & b_1 & b_2 \end{pmatrix} \tag{65}$$

and τ and τ' can be represented by :

$$\tau = \begin{pmatrix} \tau_N \\ \tau_1 \\ \tau_2 \end{pmatrix} \tag{66}$$

$$\tau' = \begin{pmatrix} \tau'_N \\ \tau'_1 \\ \tau'_2 \end{pmatrix} \tag{67}$$

With $\tau_N, \tau'_N \in \mathrm{Hom}(N, \mathcal{O}_{\mathbb{P}^3}(a))$ e $\tau_1, \tau_2, \tau'_1, \tau'_2 \in \mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}, \mathcal{O}_{\mathbb{P}^3}(a))$.
It is straightforward to check that the diagram commutes if, and only if, the following system of equations is satisfied:

$$\begin{cases} \tau'_N = \lambda\tau_N \\ a_1\tau'_1 + a_2\tau'_2 = \lambda\tau_1 \\ b_1\tau'_1 + b_2\tau'_2 = \lambda\tau_2 \end{cases} \tag{68}$$

And this happen, if, and only if, τ'_1, τ'_2 are in the same plane generated by τ_1 and τ_2 in $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a))$, that is, $\langle \tau'_1, \tau'_2 \rangle$ and $\langle \tau_1,\ \tau_2 \rangle$ are represented by the same point in $G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$, and τ'_N is a fixed multiple of τ_N. These considerations proves the following proposition:

Proposition 21. There exists a surjective map:

$$\pi : P(a,1) \to \mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)) \tag{69}$$

whose fibres are dense subsets of $\mathrm{H}^0(N(a))$, where $\mathcal{I}(2,1)$ is the moduly space of the null correlation bundles, and $N \in \mathcal{I}(2,1)$.

Proof. The map is already defined, and clearly is surjective. To see that the fibres are dense open sets of $\mathrm{H}^0(N(a))$, note that the common zero loci of τ_1 and τ_2 is a degree a^2 curve on $\mathbb{P}^3$ denoted by C, and the zero locus of τ_N is a curve of degree $2a$, so the set $\pi^{-1}(N, \langle \tau_1, \tau_2 \rangle)$ is determined by $\tau_N \in \mathrm{H}^0(N(a))$ surjective such that τ_N is non zero on C, which is dense in $\mathrm{H}^0(N(a))$. □

Now we will prove that $P(a,1)$ can be identified with an open dense subset of a vector bundle $\pi : \mathcal{P} \to \mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$ and therefore it will be a a irreducible smooth variety.

Note that each $N \in \mathcal{I}(2,1)$ can be identified with a point in $\mathbb{P}(\mathrm{H}^0(\Omega_{\mathbb{P}^3}(2)))$, since N is by definition, cokernel of an injective map $\sigma \in \mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(-1), \Omega_{\mathbb{P}^3}((1))$ modulo isomorphism, for this reason we will denote N by N_σ, and $\mathcal{I}(2,1)$ can be identified with the open dense subset $U \subset \mathbb{P}(\mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(-1), \Omega_{\mathbb{P}^3}(1))) \simeq \mathbb{P}(\mathrm{H}^0(\Omega_{\mathbb{P}^3}(2)))$ of the injective sections. We have the following exact sequence:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^3}(a-1)) \xrightarrow{\sigma} \Omega_{\mathbb{P}^3}(a+1) \longrightarrow N_\sigma(a) \longrightarrow 0 \tag{70}$$

Therefore by the long exact sequence of cohomology there exists a natural isomorphism of $\mathrm{H}^0(N_\sigma(a))$ with the quotient vector space $\mathrm{H}^0(\Omega_{\mathbb{P}^3}(a+1))/\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a-1))$.

Note that $U \subseteq \mathbb{P}(\mathrm{Hom}(\mathcal{O}_{\mathbb{P}^3}(a-1), \Omega_{\mathbb{P}^3}(a+1))) \simeq \mathbb{P}(\mathrm{H}^0(\Omega_{\mathbb{P}^3}(2)))$, define the morphism $\tilde{\sigma}$ as the multiplication of cordinates :

$$0 \longrightarrow \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a-1)) \otimes \mathcal{O}_U(-1) \xrightarrow{\tilde{\sigma}} \mathrm{H}^0(\Omega_{\mathbb{P}^3}(a+1)) \otimes \mathcal{O}_U \tag{71}$$

The cokernel of this map is again a vector bundle over U, hence over $\mathcal{I}(2,1)$ denoted by $\mathcal{P}(a)$, such that for each point $\sigma \in \mathbb{P}(\mathrm{H}^0(\Omega_{\mathbb{P}^3}(2)))$ the fibre at σ is $\mathrm{H}^0(\mathrm{coker}\,\sigma(a)) =: \mathrm{H}^0(N_\sigma(a))$. Now consider the projection on the first coordinate $p_1 : \mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a))) \to \mathcal{I}(2,1)$. The fibers at each point of the vector bundle $p_1^*\mathcal{P}(a)$ are again isomorphic to $\mathrm{H}^0(N_\sigma(a))$.

This proves that there exists a vector bundle $\pi : p_1^*\mathcal{P}(a) \to \mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$, such that $P(a,1)$ is a dense open subset of the total space of $p_1^*\mathcal{P}$, since $\pi : p_1^*\mathcal{P}(a) \to \mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$ is a vector bundle, the total space of $p_1^*\mathcal{P}(a)$ is non-singular and irreducible, then, it is possible to induce a structure of variety on $P(a,1)$, making it non-singular and irreducible. Putting all these results together we have the following Theorem.

Theorem 22. $P(a,1)$ is an open subset of the total space of the vector bundle over $\mathcal{I}(2,1) \times G(2, \mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(a)))$ whose fibres are $\mathrm{H}^0(N(a))$. Thus is an irreducible, non-singular variety of dimension $a\binom{a+3}{a+1} + 2\binom{a+3}{3} - \binom{a+2}{3} + 1$.

The dimension of the variety $P(a,1)$ given by this theorem coincides with the dimension with the dimension of the family of vector bundles $\mathcal{F}(a,1)$ given by Corollary 11. To each point $\mathbf{t} := [\tilde{E}, \tau] \in P(a,1)$ one can associate the vector bundle $E(\mathbf{t}) \in \mathcal{B}(0, a^2+1)$. We conjecture that the image of $P(a,1)$ in $\mathcal{B}(a^2+1)$ by this map is an irreducible oversized component for every $a \geq 4$. For $a = 3$, $P(3,1)$ has dimension 76 which is too small to be an irreducible component of $\mathcal{B}(10)$, since its irreducible components has dimension at least 77, and in the next section we will show that the image of $P(2,1)$ is an irreducible component of $\mathcal{B}(5)$.

7. The number of components of $\mathcal{B}(5)$

In [10] Hartshorne and Rao proved that any rank 2 stable vector bundle in $\mathcal{B}(5)$ is cohomology of one of the following monads:

$$0 \to 5\cdot\mathcal{O}_{\mathbb{P}^3}(-1) \to 12\cdot\mathcal{O}_{\mathbb{P}^3} \to 5\cdot\mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{72}$$

$$0 \to 2\cdot\mathcal{O}_{\mathbb{P}^3}(-2) \to 3\cdot\mathcal{O}_{\mathbb{P}^3}(-1) \oplus 3\cdot\mathcal{O}_{\mathbb{P}^3}(1) \to 2\cdot\mathcal{O}_{\mathbb{P}^3}(2) \to 0 \tag{73}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-3) \to \mathcal{O}_{\mathbb{P}^3}(-2) \oplus 2\cdot\mathcal{O}_{\mathbb{P}^3} \oplus \mathcal{O}_{\mathbb{P}^3}(2) \to \mathcal{O}_{\mathbb{P}^3}(3) \to 0 \tag{74}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-2) \oplus \mathcal{O}_{\mathbb{P}^3}(-1) \to 6\cdot\mathcal{O}_{\mathbb{P}^3} \to \mathcal{O}_{\mathbb{P}^3}(1) \oplus \mathcal{O}_{\mathbb{P}^3}(2) \to 0 \tag{75}$$

$$0 \to \mathcal{O}_{\mathbb{P}^3}(-2)\oplus 2\cdot\mathcal{O}_{\mathbb{P}^3}(-1) \xrightarrow{\alpha} \mathcal{O}_{\mathbb{P}^3}(-1)\oplus 6\cdot\mathcal{O}_{\mathbb{P}^3} \oplus \mathcal{O}_{\mathbb{P}^3}(1) \xrightarrow{\beta} \mathcal{O}_{\mathbb{P}^3}(2)\oplus 2\cdot\mathcal{O}_{\mathbb{P}^3}(1) \to 0 \tag{76}$$

For each stable rank two vector bundle E on $\mathbb{P}^3$ with 0 first Chern class the number $h^1(E(-2)) \mod 2$ is called the Atiyah–Rees invariant of E. Hartshorne showed in [9, Corollary 2.4] that this number is invariant on the components of the moduli space of stable vector bundles on $\mathbb{P}^3$.

The remain of this section will be devoted to prove the main Theorem of this work

Main Theorem. The moduli space $\mathcal{B}(5)$ has exactly 3 irreducible components. Namely:

i) the instanton component, of dimension 37, which contains the bundles given as cohomology of monads of the form (72) and (73)

ii) the Ein component, of dimension 40, which contains the bundles given as cohomology of monads of the form (74)

iii) the closure of $\mathcal{F}(2,1)$ in $\mathcal{B}(2,0,5)$, of dimension 37, which contains the bundles given as cohomology of monads of the form (75) and (76).

The vector bundles cohomology of monads of the form (72) and (73) have Atiyah–Rees invariant equals 0 and, Rao showed in [19] that they lie on the same component in $\mathcal{B}(5)$, whose generic vector bundle is cohomology of monads (72) and the vector bundles cohomology of the monads (73) lie in its closure. Therefore there is exactly one component of vector bundles with Atiyah–Rees invariant equal to 0 in $\mathcal{B}(5)$, and this is item i) of the Theorem 7.

Concerning the vector bundles with Atiyah–Rees invariant equal to 1, Hartshorne proved in [8] that the closure of the family of vector bundle arising as cohomology of the monads of the form (74) is an oversized component of $\mathcal{B}(5)$ of dimension 40, and this is item ii) of the Theorem 7. Because of the study of such monads made by Ein in [7] we call this family of vector bundles of generalized null correlation bundle and denote this family by $\mathcal{E}$.

It is straighfoward to check that the Atiyah–Rees invariant of every vector bundle $E \in \mathcal{F}(2,1)$ is 1, thus, in principle $\mathcal{F}(2,1)$ could be contained in the closure of $\mathcal{E}$. We will dedicate the rest of this section to show that this does not happen.

Using the notation of the Lemma 3, consider the point $\mathbf{t} \in P(2,1)$ associated with the vector bundle $E(\mathbf{t})$ which is cohomology of the monad (33), with the maps $a_0 = \alpha$ and $b_0 = \beta$, defined given by the equations (21) and (22). Fixing a basis for $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(1))$, $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(2))$, $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(3))$ and $\mathrm{H}^0(\mathcal{O}_{\mathbb{P}^3}(4)))$. One checks explicitly that d_0 is surjective, hence by Lemma 3 $\mathrm{Ext}^2(E(\mathbf{t}), E(\mathbf{t})) = \operatorname{coker} d_0 = 0$. Since $E(\mathbf{t})$ is stable, one has $\dim \mathrm{Ext}^1(E(\mathbf{t}), E(\mathbf{t})) = \dim P(2,1) = 37$. Since $P(2,1)$ is irreducible, there exists an open dense in $U \subseteq P(2,1)$, such that for each $\mathbf{t} \in U$, one has $\dim \mathrm{Ext}^1(E(\mathbf{t}), E(\mathbf{t})) = \dim P(2,1)$. This implies that, the closure of the image of $P(2,1)$ into $\mathcal{B}(5)$ is an irreducible component. Furthermore, this fact, and Proposition 20 implies that the image of $P(2,1)$ in $\mathcal{B}(5)$ coincides with $\mathcal{F}(2,1)$. Hence $\mathcal{F}(2,1)$ is contained in exactly one irreducible component. We we will this irreducible component by $\overline{\mathcal{F}(2,1)}$. We can summarize this in the next Proposition

Proposition 23. $\overline{\mathcal{F}(2,1)}$ is a generically smooth irreducible component of $\mathcal{B}(5)$ of dimension 37.

In order to give a complete description of the vector bundles $E \in \mathcal{F}(2,1)$, we include here its cohomology table that can be computed by the display of the monad and Serre Duality.

We will use the Theorem 23 to determine the number of irreducible components of $\mathcal{B}(5)$.

TABLE 1. $\dim \mathrm{H}^i(E(l))$, for a general $E \in \mathcal{F}(2,1)$

$i\backslash l$	-4	-3	-2	-1	0
3	0	0	0	0	0
2	8	5	1	0	0
1	0	0	1	5	8
0	0	0	0	0	0

Proposition 24. Consider $\mathcal{F}(2,1)$ and $\mathcal{E}$ be as defined before, then $\mathcal{F}(2,1) \cap \overline{\mathcal{E}} = \emptyset$.

Proof. By the display of the monad (74), it is possible to check that for every vector bundle $E \in \mathcal{E}$ one has $\dim \mathrm{H}^1(E(-2)) = 3$, and for every $F \in \mathcal{F}(2,1)$ one has by the Table 1 that $\dim \mathrm{H}^1(F(-2)) = 1$. Suppose by contradiction that $\mathcal{F}(2,1) \cap \overline{\mathcal{E}} \neq \emptyset$, then there exists a vector bundle in $E \in \mathcal{F}(2,1)$ such that $E \in \overline{\mathcal{E}}$. By the inferior semi-continuity of the dimension of the cohomology groups of coherent sheaves one has that $\dim \mathrm{H}^1(E(-2)) \geq 3$, but since $E \in \mathcal{F}(2,1)$ we have $\dim \mathrm{H}^1(E(-2)) = 1$, which can not occur. Thus $\mathcal{F}(2,1) \cap \overline{\mathcal{E}} = \emptyset$. □

This Proposition proves that the family of vector bundles $\mathcal{F}(2,1)$ lies in a irreducible component on the moduli space $\mathcal{B}(5)$, different from that one described by Ein. Therefore $\overline{P(2,1)}$ is a new component of $\mathcal{B}(5)$.

In order to establish the exact number of components of $\mathcal{B}(5)$, we focus on the family of vector bundles arising as cohomology of monads of the form (76). As mentioned before, the Atiyah–Rees invariant of the vector bundles in this family is equal to 1. By [15, Thm 2.5], one sees that its dimension is at most 36, thus it is too small to contain any irreducible component and therefore must lie in $\overline{\mathcal{F}(2,1)}$ or in the closure of $\mathcal{E}$. Indeed, we prove in the next proposition that the later case does not happen.

Proposition 25. The family of vector bundles cohomology of the monads of the form (76) lies in $\overline{P(2,1)}$

Proof. Suppose by contradiction that there exists a vector bundle E cohomology of a monad of the form 76) such that $E \in \overline{\mathcal{E}}$, again, by the inferior semi-continuity of the dimension of the cohomology groups of coherent sheaves one has that $\dim \mathrm{H}^1(E(-2)) \geq 3$ but by the display of the monad (76) one can check that $\dim \mathrm{H}^1(E(-2)) = 1 < 3$, hence it can not occur. Thus the family of vector bundles cohomology of monads of the form (76) lies in $P(2,1)$. □

Therefore this proposition concludes the proof of item iii) hence the proof of Theorem 7.

References

[1] C. Banica, N. Manolache, Rank 2 stable vetor bundles on $\mathbb{P}^3$, with Chern Classes $c_1 = -1, c_2 = -4$. Math. Z. **190** (1985), 315–339.

[2] W. Barth, Some properties of stable rank-2 vector bundles on $\mathbb{P}^n$. Math. Ann. **226** (1977), 125–150.

[3] W. Barth, Stable vector bundles on $\mathbb{P}^3$, some experimental data. Asterisque **71** (1980), 205–218

[4] G. Bohnhorst,H. Spindler. The stability of certain vector bundles on $\mathbb{P}^n$ (pp. 39-50). Springer Berlin Heidelberg (1992).

[5] U. Bruzzo, D. Markushevich, and A. Tikhomirov. Moduli of symplectic instanton vector bundles of higher rank on projective space $\mathbb{P}^3$. Cent. Eur. J. Math. **10** (2012): 1232–1245.

[6] L. Costa, R. M. Miró-Roig. Monads and regularity of vector bundles on projective varieties. Michigan Math. J. **55** (2007), 417–436.

[7] L. Ein, Generalized null correlation bundles, Nagoya Math. J. **111** (1988), 13–24.

[8] R. Hartshorne, Stable reflexive sheaves. Math. Ann. **254** (1980), 121–176.

[9] R. Hartshorne, Stable vector bundles of rank 2 on $\mathbb{P}^3$. Math. Ann. **238** (1978), 229–280.

[10] R. Hartshorne, A. P. Rao. Spectra and monads of stable bundles. J. Math. Kyoto Univ. **31** (1991), 789–806.

[11] R. Hartshorne, I. Sols, Stable rank 2 vector bundles on $\mathbb{P}^3$ with $c_1 = -1, c_2 = 2$. J. Reine Angew. Math. **325** (1981), 145–152.

[12] G. Horrocks, Vector bundles on the punctured spectrum of a local ring. Proc. London Math. Soc. **14** (1964), 689–713.

[13] M. Jardim, Instanton sheaves on complex projective spaces. Collect. Math. **57** (2006), 43–68.

[14] M. Jardim, S. Marchesi, A. Wissdorf, Moduli of autodual instanton bundles, Bol. Soc. Brasil. Mat. (N.S.) **47** (2016), 823–843.

[15] M. Jardim, R.V. Martins. Linear and Steiner bundles on projective varieties. Comm. Algebra **38** (2010), 2249-2270.

[16] M. Jardim, M. Verbitsky. Trihyperkahler reduction and instanton bundles on $\mathbb{CP}^3$. Compos. Math. **150** (2014), 1836–1868.

[17] M. Maruyama : Moduli of stable sheaves. I. J. Math. Kyoto Univ. **17** (1977), 91–126.

[18] C. Okonek, M. Schneider, H. Spindler, H. S. I. Gelfand, Vector bundles on complex projective spaces. Boston: Birkhauser (1980).

[19] A.P. Rao, A family of vector bundles on $\mathbb{P}^3$ (pp. 208-231). Space Curves. Springer Berlin Heidelberg, (1981).

[20] A.S Tikhomirov, Moduli of mathematical instanton vector bundles with even on projective space. Izvestiya: Mathematics **77** (2013), 1195.

[21] A.S Tikhomirov, "Moduli of mathematical instanton vector bundles with odd on projective space." Izvestiya: Mathematics **76** (2012), 991.